

Electronic mosquito repellents for preventing mosquito bites and malaria infection (Review)

Enayati AA, Hemingway J, Garner P



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2008, Issue 1

<http://www.thecochranelibrary.com>



TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	1
BACKGROUND	2
OBJECTIVES	2
CRITERIA FOR CONSIDERING STUDIES FOR THIS REVIEW	2
SEARCH METHODS FOR IDENTIFICATION OF STUDIES	3
METHODS OF THE REVIEW	3
DESCRIPTION OF STUDIES	3
METHODOLOGICAL QUALITY	4
RESULTS	4
DISCUSSION	4
AUTHORS' CONCLUSIONS	5
POTENTIAL CONFLICT OF INTEREST	5
ACKNOWLEDGEMENTS	5
SOURCES OF SUPPORT	5
REFERENCES	5
TABLES	6
Characteristics of included studies	6
Characteristics of excluded studies	10
ADDITIONAL TABLES	11
Table 01. Detailed search strategies	11
Table 02. Study methodological quality	11
Table 03. Mosquito landing rates	12
GRAPHS AND OTHER TABLES	12
INDEX TERMS	12
COVER SHEET	13

Electronic mosquito repellents for preventing mosquito bites and malaria infection (Review)

Enayati AA, Hemingway J, Garner P

This record should be cited as:

Enayati AA, Hemingway J, Garner P. Electronic mosquito repellents for preventing mosquito bites and malaria infection. *Cochrane Database of Systematic Reviews* 2007, Issue 2. Art. No.: CD005434. DOI: 10.1002/14651858.CD005434.pub2.

This version first published online: 18 April 2007 in Issue 2, 2007.

Date of most recent substantive amendment: 16 February 2007

ABSTRACT

Background

Electronic mosquito repellents (EMRs) are marketed to prevent mosquitoes biting and to prevent malaria.

Objectives

To assess whether EMRs prevent mosquito bites, and to assess any evidence of an effect on malaria infection.

Search strategy

In August 2006, we searched the Cochrane Infectious Diseases Group Specialized Register, CENTRAL, MEDLINE, EMBASE, LILACS, Cambridge Scientific Abstracts, and the Science Citation Index. We also checked conference proceedings, contacted international specialist centres and EMR manufacturers, and checked reference lists.

Selection criteria

Field entomological studies, which controlled for geographic site, time, and attractiveness of human participants, of EMRs for preventing mosquito bites; and randomized and quasi-randomized controlled trials of EMRs to prevent malaria infection.

Data collection and analysis

Two authors assessed trial quality, and extracted and analysed the data.

Main results

Ten field entomological studies met the inclusion criteria. All 10 studies found that there was no difference in the number of mosquitoes caught from the bare body parts of the human participants with or without an EMR. No randomized or quasi-randomized controlled trials on the efficacy of EMR on malaria infection were found.

Authors' conclusions

Field entomological studies confirm that EMRs have no effect on preventing mosquito bites. Therefore there is no justification for marketing them to prevent malaria infection.

PLAIN LANGUAGE SUMMARY

Electronic mosquito repellents for preventing mosquito bites and malaria infection

Malaria is a major health problem that particularly affects people living in sub-Saharan Africa and other tropical parts of the world. It often causes considerable morbidity and mortality especially in children under five. It is transmitted by mosquito bites from infected female mosquitoes. Several strategies and approaches are available for preventing mosquito bites and malaria infection, including repellents, and these approaches will be considered by those living in affected areas and by travellers to areas where there is high risk of infection. Electronic mosquito repellents (EMRs) are designed to repel female mosquitoes by emitting high-pitched sounds almost inaudible to the human ear. EMRs are claimed by their manufacturers to be effective in repelling mosquitoes and preventing disease.

No randomized controlled trials were found, but 10 field studies looking at the number of mosquitoes caught on the bare body parts of humans were assessed. These studies were conducted in various parts of the world with different species of mosquitoes and were controlled for factors such as locality and timing. One study used just one observer with seven observations, while the highest assessment included 18 observers with 324 observations. There was no evidence in the field studies to support any repelling effects of EMRs, hence no evidence to support their promotion or use. Future randomized controlled trials are not proposed as there was no suggestion in the field studies that EMRs show any promise as a preventive measure against malaria.

BACKGROUND

Malaria affects more than 250 million people and causes more than a million deaths each year (WHO 2005). One important control strategy against this and other mosquito-borne diseases is mosquito control, which aims to reduce human-mosquito contact. Different control measures are used routinely against mosquitoes and their larvae, including chemical (eg insecticide), biological (eg larvivorous fish or pathogenic fungi), environmental (eg land filling or drainage), and personal protection (eg mosquito repellents formulated as pills, coils, ointments, lotions, and sprays; and insecticide-treated or untreated bed nets).

Electronic mosquito repellents (EMRs) are marketed in response to a huge demand from the public for convenient, safe, and effective antimosquito products. Female *Anopheles* mosquitoes transmit malaria by sucking blood from humans, and these small hand-held, battery-powered EMRs are intended to repel them by emitting a high frequency buzz almost inaudible to the human ear. They can be used both indoors and outdoors, and are claimed to repel mosquitoes within a range of up to 2.5 metres (Kutz 1974; Helson 1977). No adverse effects have been reported in the literature. Mobile phone companies also market a ring tone that is claimed to repel mosquitoes within a one-metre radius (BBC 2003).

Some of the EMRs seem to be based on known aspects of mosquito behaviour, while others have no scientific data to substantiate their claims. Manufacturers have put forward at least two reasons to explain the alleged repellent action of sound against mosquitoes. One reason is that the flight sound of males repels females once they have been inseminated (Foster 1985); hence, whatever mimics the males' flight sound may repel females. However, research has shown that male mosquitoes are actually the ones attracted by the female flight sound and females normally have a very weak sensitivity for sound compared with the males (Wigglesworth 1965; Chapman 1982; McIver 1985; Michelsen 1985). Another reason is that mosquitoes avoid the ultrasonic cries of bats (Foster 1985). Although both explanations may be conceivable, there is no published scientific information to support either idea.

Different brands of EMRs have been examined for their efficacy under laboratory conditions, none of which showed any effects for the devices tested (Singleton 1977; Curtis 1982; Iglisch 1983; Foster 1985; Jensen 2000; Andrade 2001; Cabrini 2006). There

are review articles concluding that the EMRs are ineffective in repelling mosquitoes (Coro 1998; Coro 2000). Scientific skepticism over the last 30 years and a successful prosecution of EMR sellers under the UK Trade Description Act in 1980s (Curtis 1994; BBC 2005) seems to have done little to deter manufacturers marketing EMRs and the people who buy them. This is a concern because it is likely to lead to consumers not using other protective methods that are proven to work. This could result in an increased risk of infection with mosquito-borne diseases, especially malaria (Jensen 2000).

Despite the scientific view and research findings, EMRs are still widely promoted and used by the public. We therefore decided to systematically review all reliable research about the effects of high-pitched sounds in preventing mosquito bites and, hence, to assess whether there is any evidence that EMRs have any potential in preventing malaria in the field setting. We included only field studies since laboratory studies do not reflect influences on mosquito behaviour, including climate, mosquito density, and composition of different species in the same locality.

OBJECTIVES

To assess whether EMRs prevent mosquito bites, and to assess evidence of impact on malaria infection.

CRITERIA FOR CONSIDERING STUDIES FOR THIS REVIEW

Types of studies

Preventing mosquito bites

Field entomological studies that control for geographic site (conducted in same locality), time (conducted at same time), and attractiveness of human participants (by rotating participants between the experiments with and without the EMR).

Preventing malaria infection

Randomized and quasi-randomized controlled trials.

Types of participants

Adults or children.

Types of intervention

Intervention

EMRs with any operational wavelength.

Control

Dummy EMRs, inoperable EMRs, EMRs switched off, or no EMRs. We excluded other repellents and treated or untreated bed nets as controls.

If used, malaria chemoprophylaxis must be identical in both the intervention and control groups.

Types of outcome measures

- The number of mosquitoes of any species landing on exposed body parts of humans acting as baits. Time period defined by entomological collection procedures.
- Malaria infection, defined as clinical malaria (fever with malaria parasitaemia detected by microscopy or rapid test); or asymptomatic malaria parasitaemia.

SEARCH METHODS FOR IDENTIFICATION OF STUDIES

See: Cochrane Infectious Diseases Group methods used in reviews.

We attempted to locate all relevant studies regardless of language or publication status (published, unpublished, in press, and in progress).

Databases

The following databases were searched using the search terms and strategy described in Table 01: Cochrane Infectious Diseases Group Specialized Register (August 2006); Cochrane Central Register of Controlled Trials (CENTRAL), published in *The Cochrane Library* (2006, Issue 2); MEDLINE (1966 to August 2006); EMBASE (1974 to August 2006); LILACS (1982 to August 2006); Cambridge Scientific Abstracts (CSA) (1982 to August 2006); and Science Citation Index (SCI) (1945 to August 2006).

Conference proceedings

The following conference proceedings were searched for relevant abstracts: XV International Congress of Tropical Medicine and Malaria, Cartagena, Colombia, August 2000; First MIM Pan-African Malaria Conference, Dakar, Senegal, 6 to 9 January 1997; Second MIM Pan-African Malaria Conference, Durban, South Africa, 15 to 19 March 1999; Third MIM Pan-African Malaria Conference, Arusha, Tanzania, 17 to 22 November 2002; Fourth MIM Pan-African Malaria Conference, Yaoundé, Cameroon, 13 to 18 November 2005; International Conference on Entomology, Brisbane, Australia, 15 to 21 August 2004; and Medicine and Health in the Tropics, Marseille, France, 11 to 15 September 2005.

Researchers, organizations, and manufacturers

We contacted some corresponding authors and field and clinical experts (Professor Chris Curtis, London School of Hygiene and Tropical Medicine; Dr Morteza Zaim, WHO Pesticide Evaluation Scheme (WHOPES), Geneva, to enquire about other published or unpublished relevant studies (September 2006). We also contacted EMR manufacturers (Isotronic, Lentek International Inc., Electronic Pest Controls Ltd.) for unpublished and ongoing trials or studies (September 2005).

Reference lists

We also checked the reference lists of all studies identified by the above methods.

METHODS OF THE REVIEW

AAE scanned the results of the literature search for potentially relevant studies and then retrieved the full articles. AAE and PG independently assessed the potentially relevant studies using an eligibility form based on the inclusion criteria; disagreements were resolved through discussion.

We independently extracted data from all included studies using a data extraction form and resolved any disagreements in the extracted data by referring to the original paper and through discussion. We described the devices tested and the number of observations made, and assessed the quality of the studies in relation to whether they controlled for study locality, time of day or night, participants (used same people to bait mosquitoes), and whether the observers were blinded. We also assessed the number of times observations were repeated to gain some quantitative measure of quality and grouped this with an arbitrary cut off into adequate (20 or more) or inadequate (less than 20). We summarized the results of the methodological quality assessment in Table 02.

The number of landings on which the rates were calculated varied considerably according to different ecological and geographical situations, mosquito species, and season and time of day of the tests. If possible, we would have tested for a difference using the original data to measure a mean difference between arms within one study and to calculate 95% confidence intervals. Had a difference been shown, we would also have examined the effects of EMR by a variety of factors: EMR frequency (< and \geq 20 kHz); mosquito population density; malaria endemicity (< and \geq entomological inoculation rate of 1/person/night); and mosquito species.

DESCRIPTION OF STUDIES

We identified 18 potentially relevant studies of EMR to prevent mosquito bites and included 10 (see 'Characteristics of included

studies' for study details); none were randomized or quasi-randomized controlled trials that used EMR to prevent malaria. We excluded eight studies because they were only laboratory based or because they did not provide any data or did not control for locality, time, and blinding (see 'Characteristics of excluded studies'). The papers for the 10 included studies contained 22 experiments, of which 15 were field experiments that met the reviews inclusion criteria; the excluded seven experiments were only laboratory based or used chemical repellents.

Seven studies were carried out in the North America, three in Canada (Helson 1977; Belton 1981; Lewis 1982) and four in the USA (Gorham 1974; Kutz 1974; Garcia 1976; Schreck 1977). Two studies were done in Africa, in Gabon (Sylla 2000) and in The Gambia (Snow 1977). One study was undertaken in Russia (Rasnitsyn 1974).

Seven studies gave the commercial name of the EMRs tested (Rasnitsyn 1974; Helson 1977; Schreck 1977; Snow 1977; Belton 1981; Lewis 1982; Sylla 2000); five studies gave some information about the ultrasound frequencies used, which ranged from 125 Hz to 74,600 Hz (Kutz 1974; Belton 1981; Rasnitsyn 1974; Snow 1977; Sylla 2000). The other studies gave no commercial name and no details of the frequencies used.

All studies counted mosquitoes landing on the bare body parts (mostly arms, legs and/or feet) of the human participants for definite time periods with the EMR switched on or off, or, in some studies, with or without a functional EMR as case and control. None of the field studies performed in North America and Russia were on *Anopheles* mosquitoes; they used *Aedes*, *Culex*, *Culiseta*, and *Mansonia* mosquitoes. The two studies in Africa were on *Anopheles* as well as other mosquitoes. The lowest number of observers was one (with 7 observations; Gorham 1974) and the highest was 18 (with 324 observations; Sylla 2000). Also, timing and length of collections varied, ranging from one minute (Kutz 1974) to over a 12-hour period (Sylla 2000).

METHODOLOGICAL QUALITY

As shown in Table 02, all studies made some attempts to control for locality (geography) of the study area (wooded area, forest, plain, beach) as they measured landing rates with and without the EMR in the same geographical area. Different individuals may have different attraction for the mosquitoes, and the studies controlled for this, usually by swapping the EMR between the participants so that the same individuals acted as both case and control.

The density of mosquitoes and the intensity with which they attempt to bite varies throughout day and night. Our inclusion criteria required studies to attempt to control for time. This was clearly described in all but one study (Garcia 1976).

One study blinded the observers to whether they were measuring during a control or experimental phase (Sylla 2000). The use of blinding was unclear or not used in the other studies.

In order to test significance, we intended to consider rates per person in participants in control or intervention areas. As landing rates were not given per person, we were unable to compare these data statistically. We recorded the number of times observations were repeated to gain some insight into data quality, arbitrarily defining this as adequate (with 20 or more repeated observations or human participants) and inadequate (less than 20 observations). Three studies were of adequate quality by this criterion.

RESULTS

The number of mosquitoes landing per collection with and without EMR are presented in Table 03. All 10 studies reported that the landing rates with and without the EMR were little different and that the EMRs failed to repel mosquitoes. These results occurred regardless of the study location, mosquito density, mosquito genera, or time of study (ie day or night with day-biting and night-biting mosquitoes).

No trials were found to assess the effects of EMRs on malaria infection.

DISCUSSION

The included studies were of good quality, had controlled for locality, and all but one had explicitly controlled for time of day or night, and the human bait for the mosquitoes. The results of this review provide clear evidence from field-based studies that there is no hint that these devices have any effect on mosquito landing rates. The studies reported here examined the effectiveness of the EMRs with different methods, settings, mosquito species (since they may respond differently to the high-pitched sounds emitted by the EMRs), frequencies of the sound emitted by the EMRs (since mosquitoes may respond to a particular sound wavelength), and times of day (since day-biting and night-biting mosquitoes may behave differently to the sound emitted by the EMRs), and mosquito density (since this may affect EMR efficacy), but none of them supported the claims of the EMRs' effectiveness. Although we did not conduct a meta-analysis of the included studies, there was no suggestion of difference in landing rates between cases and controls in any trial. In 12 of the 15 experiments, the landing rates in the groups with functioning EMR was actually higher than in the control groups. The absolute number of mosquitoes landing on the human participants during the experiments while the EMR was functioning was too high to consider the EMR a repellent.

EMRs are claimed to be effective by mimicking the sound waves produced by the beating of male mosquitoes' wings, especially during swarms. Female mosquitoes, which bite humans, are claimed

to be repelled by this sound since they mate only once in their lives. Hence the repellent mechanism should be based on the hearing mechanism in females. However, this theory is implausible since the hearing ability of the females is relatively weak (Wigglesworth 1965; Chapman 1982; Michelsen 1985). It is the hearing system of males that is relatively strong, and the presence of numerous sound and vibration receptors (known as Johnston organ) on their plumose antennae enables them to detect the vibration in the environment as well as the sound of female mosquitoes (Chapman 1982). Thus it is not surprising that the included studies did not produce any evidence that EMRs act as repellents.

AUTHORS' CONCLUSIONS

Implications for practice

EMRs are not effective in repelling mosquitoes and should not be recommended or used.

Implications for research

There is no evidence of an effect of EMRs on landing rates. Thus there is no evidence that these EMRs could potentially be useful in preventing malaria in humans. Given these findings from 10 carefully conducted studies, it would not be worthwhile to conduct further research on EMRs in preventing mosquitoes biting or in trying to prevent the acquisition of malaria.

POTENTIAL CONFLICT OF INTEREST

None known.

ACKNOWLEDGEMENTS

AA Enayati developed the protocol of the review during the Fellowship Programme organized by the Cochrane Infectious Diseases Group in March 2005. The Department for International Development (DFID) UK supports this Programme through the Effective Health Care Research Programme Consortium at the Liverpool School of Tropical Medicine.

This document is an output from a project funded by the DFID for the benefit of developing countries. The views expressed are not necessarily those of DFID.

SOURCES OF SUPPORT

External sources of support

- Department of International Development UK

Internal sources of support

- Liverpool School of Tropical Medicine UK

REFERENCES

References to studies included in this review

- Belton 1981** {published data only}
Belton P. An acoustic evaluation of electronic mosquito repellers. *Mosquito News* 1981;**41**:751–5.
- Garcia 1976** {published data only}
Garcia R, Des Rochers B, Voigt WG. Evaluation of electronic mosquito repellers under laboratory and field conditions. *Vector Views* 1976;**23**(5/6):21–3.
- Gorham 1974** {published data only}
Gorham JR. Tests of mosquito repellents in Alaska. *Mosquito News* 1974;**34**:409–15.
- Helson 1977** {published data only}
Helson BV, Wright RE. Field evaluation of electronic mosquito repellers in Ontario. Proceedings of the Entomological Society of Ontario. 1977; Vol. 108:59–61.
- Kutz 1974** {published data only}
Kutz FW. Evaluation of an electronic mosquito repelling device. *Mosquito News* 1974;**34**:369–75.
- Lewis 1982** {published data only}
Lewis DJ, Fairchild WL, Leprince DJ. Evaluation of an electronic mosquito repeller. *Canadian Entomologist* 1982;**114**(8):699–702.

- Rasnitsyn 1974** {published data only}
Rasnitsyn SP, Alekseev AN, Gornostaeva RM, Kupriianova ES, Potapov AA. Negative results of test of sound generator devices designed for mosquito repellence. *Meditsinskaya Parazitologiya i Parazitarnye Bolezni* 1974;**43**(6):706–8.
- Schreck 1977** {published data only}
Schreck CE, Weidhaas DE, Smith N. Evaluation of electronic sound-producing buzzers against *Aedes taeniorhynchus* and *Ae. sollicitans*. *Mosquito News* 1977;**37**(3):529–31.
- Snow 1977** {published data only}
Snow WF. Trials with an electronic mosquito-repelling device in West Africa. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1977;**71**(5):449–50.
- Sylla 2000** {published data only}
Sylla el-H K, Lell B, Kremsner PG. A blinded, controlled trial of an ultrasound device as mosquito repellent. *Wiener Klinische Wochenschrift* 2000;**112**(10):448–50.

References to studies excluded from this review

- Andrade 2001**
Andrade CFS, Bueno VS. Evaluation of electronic mosquito-repelling devices using *Aedes albopictus* (Skuse) (Diptera: Culicidae). *Neotropical Entomology* 2001;**30**(3):497–9.

- Arevad 1982**
Arevad K. Evaluation of an electronic mosquito-repelling device. Danish Pest Infestation Laboratory Annual Report 1982:21.
- Cabrini 2006**
Cabrini I, Andrade CFS. Evaluation of seven new electronic mosquito repellents. *Entomologia Experimentalis et Applicata* 2006;**121**(2):185–8.
- Curtis 1982**
Curtis C, White G. Once bitten, twice shy. *New Scientist* 1982;**93**: 328.
- Foster 1985**
Foster WA, Lutes KI. Tests of ultrasonic emissions on mosquito attraction to hosts in a flight chamber. *Journal of the American Mosquito Control Association* 1985;**1**(2):199–202.
- Iglisch 1983**
Iglisch VI. Mosquito-repellent efficacy of sound-wave-emitting apparatus [Zur stechmückenabweisenden Wirksamkeit von schallwellenerzeugenden Geräten]. *Anzeiger für Schadlingskde, Pflanzenschutz, Umweltschutz* 1983;**56**:135–40.
- Jensen 2000**
Jensen T, Lampman R, Slamecka MC, Novak RJ. Field efficacy of commercial antimosquito products in Illinois. *Journal of the American Mosquito Control Association* 2000;**16**(2):148–52.
- Singleton 1977**
Singleton RE. Evaluation of two mosquito-repelling devices. *Mosquito News* 1977;**37**(2):195–9.
- Additional references**
- BBC 2003**
S Korean mobiles to zap mosquitoes. news.bbc.co.uk/1/hi/world/asia-pacific/3063401.stm 14 July 2003.
- BBC 2005**
Mosquito buzzers' worth queried. news.bbc.co.uk/1/hi/health/4218234.stm 6 September 2005.
- Chapman 1982**
Chapman RF. *The insects: structure and function*. 3rd Edition. London: Hodder & Stoughton, 1982.
- Coro 1998**
Coro F, Suarez S. Electronic repellents against mosquitoes: the propaganda and the reality [Repelentes electronicos contra mosquitos: propaganda y realidad]. *Revista Cubana de Medicina Tropical* 1998;**50**(2):89–92.
- Coro 2000**
Coro F, Suarez S. Review and history of electronic mosquito repellents. *Wing Beats* 2000;**11**(2):6–7.
- Curtis 1994**
Curtis CF. Anti-mosquito buzzers, advertising and the law. *Wing Beats* 1994;**5**:10–2.
- McIver 1985**
McIver SB. Mechanoreception. In: KerkutGA, GilbertLI editor(s). *Comprehensive insect physiology, biochemistry and pharmacology*. Vol. 6, Oxford: Pergamon, 1985:71–133.
- Michelsen 1985**
Michelsen A, Larsen ON. Hearing and sound. In: KerkutGA, GilbertLI editor(s). *Comprehensive insect physiology, biochemistry and pharmacology*. 6th Edition. Oxford: Pergamon, 1985:495–556.
- WHO 2005**
Global Partnership to Roll Back Malaria. *World malaria report: 2005*. Geneva: World Health Organization, 2005.
- Wigglesworth 1965**
Wigglesworth VB. *The principles of insect physiology*. 6th Edition. London: Methuen, 1965.

T A B L E S

Characteristics of included studies

Study	Belton 1981
Methods	Design: field study of experimental intervention versus control Two human participants with an exposed arm carried out 6 x 5 min catches once with the EMR switched on and once off Number of observers: 12 Number of observations: 12
Participants	Human adults
Interventions	1. EMR switched on 2. EMR switched off (control) EMR suppliers: Murray Distributors Ltd., North Vancouver B.C.; Peak Distributions Ltd., Surrey; Moziquit supplied by Electronic pest control Ltd., Montreal P.Q. EMR frequency: 2 to 5 KHz

Characteristics of included studies (Continued)

Outcomes	Aedes and Culex mosquito landing rates/min/participant
Notes	Location: Canada The raw number of landings with EMR switched off was not provided, but it was mentioned that the differences of the landing rates with a EMR on and off was not significant
Allocation concealment	D – Not used

Study Garcia 1976

Methods	Design: field study of experimental intervention versus control Catches carried out in 18 different locations in a forest at 2, 5, and 10 min (a total period of 102 min) intervals in the late afternoon to early evening with EMR on or off Number of observers: 6 Number of observations: 30
Participants	Human adults
Interventions	1. EMR switched on 2. EMR switched off (control) EMR supplier: unclear EMR frequency: not mentioned
Outcomes	Anopheles, Aedes, and Culex mosquito landing rates/person
Notes	Location: USA
Allocation concealment	D – Not used

Study Gorham 1974

Methods	Design: field study of experimental intervention versus control 5 min landing counts carried out on 7 occasions by single catchers with and without EMR Number of observers: 1 Number of observations: 7
Participants	Human adults
Interventions	1. EMR 2. No EMR (control) EMR supplier: not mentioned EMR frequency: not mentioned
Outcomes	Aedes and Culiseta mosquito landing rates/person/h
Notes	Location: Alaska, USA
Allocation concealment	D – Not used

Study Helson 1977

Methods	Design: field study of experimental intervention versus control Participants were 30 m apart in a wooded area caught mosquitoes landing on their arm and hand for 4 min with or without 2 types of EMR. So for each, EMR 8 min collection with EMR and 8 min collection without EMR Number of observers: 5 Number of observations: 12
Participants	Human adults

Characteristics of included studies (Continued)

Interventions	1. 2 types of EMR 2. No EMR (control) EMR source: ME Electronic Mosquito Repellent; Buzz-Off repellent EMR frequency: not mentioned
Outcomes	Aedes mosquito mean landing rate/person
Notes	Location: Canada
Allocation concealment	D – Not used

Study Kutz 1974

Methods	Design: field study of experimental intervention versus control 5 x 1 min biting counts made by 2 participants walking 20 paces along a path once with the EMR on and the other time off Number of observers: 2 Number of observations: 10
Participants	Human adults
Interventions	1. EMR switched on 2. EMR switched off (control) EMR supplier: not mentioned EMR frequency: 5.2 KHz
Outcomes	Aedes and Culex mosquito mean landing rates/person/min
Notes	Location: USA
Allocation concealment	D – Not used

Study Lewis 1982

Methods	Design: field study of experimental intervention versus control 3 participants carried out catches in 3 sites 10 m apart in 4 x 45 min catching experiments each consisting of 3 x 15 min and each of those to 3 x 5-min intervals once with EMR and once without. Participants with and without EMR were rotated Number of observers: 3 Number of observations: 12
Participants	Human adults
Interventions	1. EMR 2. No EMR (control) EMR supplier: Antipic EMR frequency: not mentioned
Outcomes	Aedes and Mansonia total bites
Notes	Location: Canada
Allocation concealment	D – Not used

Study Rasnitsyn 1974

Methods	Design: field study of experimental intervention versus control 3 participants carried 15 observations x 5 min with the EMR on or off Number of observers: 3
---------	--

Characteristics of included studies (Continued)

	Number of observations: 15
Participants	Human adults
Interventions	1. EMR switched on 2. EMR switched off EMR supplier: Skeeter Skat, Anti-parasite block, local Russian produced device EMR frequency: 125 to 74,600 Hz
Outcomes	Aedes, Culiseta, and Culex mosquito landing rates/person/h
Notes	Location: Russia
Allocation concealment	D – Not used

Study Schreck 1977

Methods	Design: field study of experimental intervention versus control 4 participants carried out the catches once without the EMR and once 2 participants carried one type of EMR and the other 2 two carried other type while walking in a wooded area where Aedes taeniorhynchus mosquitoes were abundant. They stopped every 20 paces and counted the mosquitoes on their arms. After each count the mosquitoes were killed to avoid recounting Number of observers: 4 Number of observations: 44
Participants	Human adults
Interventions	1. 2 types of EMR 2. No EMR (control) EMR supplier: Buzz-Off; Norris Electronic Mosquito Repeller EMR frequency: not mentioned
Outcomes	Aedes mosquito landing rate/person
Notes	Location: USA
Allocation concealment	D – Not used

Study Snow 1977

Methods	Design: field study of experimental intervention versus control Experiment 1: 2 teams of 2 catchers each seated 30 m apart. Each catcher collected mosquitoes for 30 minutes at 2 periods during the night. In the first period, the first team were exposed to the EMR and in the second period, the second team was exposed. Repeated on 2 consecutive nights Experiment 2: single catchers (1 with EMR and other without) sat 25 m apart and made 2 x 30 min catches over 6 nights, so a total of 2 x 2 x 30 x 6 min catches. EMRs were swapped each night between catchers Experiment 3: single catchers (1 with EMR and other without) sat 25 m apart made 2 x 30 min catches over 4 nights, so a total of 2 x 2 x 30 x 4 min catches. EMRs were swapped each night between catchers Number of observers: 4 (experiment 1); 1 (experiments 2 and 3) Number of observations: 16 (experiment 1); 12 (experiment 2 and 3)
Participants	Human adults
Interventions	1. EMR 2. No EMR (control) EMR supplier: Moziquit from Electronic Pest Controls Ltd. EMR frequency: 2.2.5 KHz

Outcomes	Number of Anopheles, Aedes, and Culex mosquitoes landing
Notes	Location: The Gambia
Allocation concealment	D – Not used

Study	Sylla 2000
Methods	<p>Design: field study of experimental intervention versus control</p> <p>18 houses chosen and grouped into 9 pairs. Each pair with a pair of working and not working EMRs that were swapped every night. 18 catches were undertaken in each house, 9 with and 9 without EMR. So a total of 9 x 9 catches in 18 houses, all together 324 catches each for 12 hours were made which makes a total catch of 3888 person hours</p> <p>Number of observers: 18 Number of observations: 324</p>
Participants	Human adults
Interventions	<p>1. Functional EMRs 2. Inoperable EMRs (control)</p> <p>EMR supplier: Isotronic, Horb, Germany EMR frequency: 3 to 11 KHz</p>
Outcomes	Anopheles, Aedes, Culex, and Mansonia mosquito landing/house/night
Notes	Location: Gabon
Allocation concealment	D – Not used
Allocation concealment: not measured for these study designs; EMR: electronic mosquito repellent	

Characteristics of excluded studies

Study	Reason for exclusion
Andrade 2001	Laboratory-based EMR study
Arevad 1982	Field-based EMR, but unclear if time, locality and bait were controlled
Cabrini 2006	Laboratory-based EMR study
Curtis 1982	Laboratory-based EMR study
Foster 1985	Laboratory-based EMR study
Iglisch 1983	Laboratory-based EMR study
Jensen 2000	Field trial but no proper control used; number of caught mosquitoes pooled together for 2 chemical repellents and EMR such that mosquito landing rates with and without EMR could not be calculated
Singleton 1977	Laboratory-based EMR study
EMR: electronic mosquito repellent	

ADDITIONAL TABLES

Table 01. Detailed search strategies

Search set	CIDG SR [^]	CENTRAL	MEDLINE	EMBASE	LILACS, CAS, SCI
1	mosquito repellent	mosquito*	mosquito*	mosquito\$	mosquito
2	mosquito control	repel*	repel*	repel\$	repel*
3	--	control	control	control	control
4	--	2 or 3	2 or 3	2 or 3	2 or 3
5	--	1 and 4	1 and 4	1 and 4	1 and 4
6	--	MOSQUITO CONTROL	MOSQUITO CONTROL	ultrasound	ultrasound
7	--	5 or 6	INSECT REPELLENTS	electronic	electronic
8	--	electronic	INSECT BITES AND STINGS/PREVENTION AND CONTROL	device\$	6 or 7
9	--	device\$	5 or 6 or 7 or 8	6 or 7 or 8	5 and 8
10	--	ultrasound	ultrasound	5 and 9	--
11	--	8 or 9 or 10	electronic	--	--
12	--	7 and 11	device\$	--	--
13	--	--	10 or 11 or 12	--	--
14	--	--	9 and 13	--	--
	[^] Cochrane Infectious Diseases Group Specialized Register	Upper case: MeSH or EMTREE heading; lower case: free text term			

Table 02. Study methodological quality

Study	Locality [^]	Time [^]	Bait [^]	Observers blind	No. observations
Belton 1981	Yes	Yes	Yes	Unclear	Inadequate
Garcia 1976	Yes	Unclear	Unclear	Unclear	Adequate
Gorham 1984	Yes	Yes	Yes	Not blinded	Inadequate
Helson 1977	Yes	Yes	Yes	Unclear	Inadequate
Kutz 1974	Yes	Yes	Yes	Unclear	Inadequate
Lewis 1982	Yes	Yes	Yes	Not blinded	Inadequate
Rasnitsyn 1974	Yes	Yes	Yes	Not blinded	Inadequate
Schreck 1977	Yes	Yes	Yes	Not blinded	Adequate
Snow 1997	Yes	Yes	Yes	Unclear	Inadequate

Table 02. Study methodological quality (Continued)

Study	Locality [^]	Time [^]	Bait [^]	Observers blind	No. observations
Sylla 2000	Yes	Yes	Yes	Blinded	Adequate
[^] Controlled					

Table 03. Mosquito landing rates

Study [^]	LR with EMR ^{^^}	LR without EMR
Belton 1981	*	*
Garcia 1976	240	193
Gorham 1984	213%** greater than the control	68.78
Helson 1977: ME Electronic Mosquito repellent	8.9	8.4
Helson 1977: Buzz-Off repellent	16.8	14.9
Kutz 1974: Person A	12.2	12.8
Kutz 1974: Person B	16.8	14.2
Lewis 1982	561.5	538
Rasnitsyn 1974	500	497
Schreck 1977: Buzz-Off repellent	25	30
Schreck 1977: Norris Device	36	As above
Snow 1997: Experiment 1	157	140
Snow 1997: Experiment 2	472	425
Snow 1997: Experiment 3	177	155
Sylla 2000	23.4	22.7

LR: landing rate

[^]See 'Characteristics of included studies' for study details

[^]Using crude units, data given in paper

*Landing rates between 2 and 7/min with no significant difference between groups

**Based on recalculation of the crude data in the paper

GRAPHS AND OTHER TABLES

This review has no analyses.

INDEX TERMS

Medical Subject Headings (MeSH)

*Anopheles; Electronics; Insect Bites and Stings [*prevention & control]; Malaria [*prevention & control]; Mosquito Control [*instrumentation]; Population Density

MeSH check words
Animals; Female; Humans

COVER SHEET

Title	Electronic mosquito repellents for preventing mosquito bites and malaria infection
Authors	Enayati AA, Hemingway J, Garner P
Contribution of author(s)	AA Enayati developed and wrote the review. P Garner helped develop the review, developed and applied inclusion and quality criteria, extracted data, and helped write the review. J Hemingway initiated the review and helped with technical issues. AA Enayati is the guarantor.
Issue protocol first published	2005/3
Review first published	2007/2
Date of most recent amendment	20 February 2007
Date of most recent SUBSTANTIVE amendment	16 February 2007
What's New	2007, Issue 2: Review first published.
Date new studies sought but none found	Information not supplied by author
Date new studies found but not yet included/excluded	Information not supplied by author
Date new studies found and included/excluded	Information not supplied by author
Date authors' conclusions section amended	Information not supplied by author
Contact address	Dr Ahmadali Enayati Lecturer Medical Entomology, School of Public Health and Environmental Health Research Centre Mazandaran University of Medical Sciences KM 18 Khazarabad Road Sari Mazandaran 48175-1553 IRAN E-mail: tmaae@liverpool.ac.uk; ahmadali_enayati@yahoo.com Tel: +98 151 3268855 Fax: +98 151 3542473
DOI	10.1002/14651858.CD005434.pub2
Cochrane Library number	CD005434
Editorial group	Cochrane Infectious Diseases Group
Editorial group code	HM-INFECTN